

Chap 5: Fourier series for periodic functions

Let $f : (-\pi, \pi] \rightarrow \mathbb{R}$ be a bounded piecewise continuous function which we continue to be a 2π -periodic function defined on \mathbb{R} , i.e.

$$f(x + 2\pi) = f(x), \quad \forall x \in \mathbb{R}.$$

The Fourier series of this function is written as

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx))$$

where

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx \quad \text{and} \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx.$$

Sufficient conditions for

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx))$$



will be considered later.

Examples of 2π periodic functions

Let $f_1(x)$ and $f_2(x)$ be defined on $(-\pi, \pi]$ as follows.

$$f_1(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq \pi, \\ 0, & \text{if } -\pi < x < 0, \end{cases} \quad \text{and} \quad f_2(x) = |x|.$$

Both of these are continued 2π -periodically and a sketch of both is shown below.



Functions which are orthogonal on $(-\pi, \pi)$

Functions f and g defined on $(-\pi, \pi)$ are orthogonal if

$$\int_{-\pi}^{\pi} f(x)g(x) dx = 0.$$

The following functions are orthogonal to each other in this sense.

$$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots, \cos(nx), \sin(nx), \dots$$

These functions are also 2π -periodic with 2π being the least period for $\cos(x)$ and $\sin(x)$.

Addition formulas to help evaluate the integrals

In complex form we have

$$e^{i(a+b)} = e^{ia}e^{ib}, \quad e^{i(a-b)} = e^{ia}e^{-ib}.$$

By adding and subtracting the expanded version of the above and taking real and imaginary parts leads to

$$\cos(a)\cos(b) = \frac{\cos(a+b) + \cos(a-b)}{2},$$

$$\sin(a)\cos(b) = \frac{\sin(a+b) + \sin(a-b)}{2},$$

$$\sin(a)\sin(b) = \frac{\cos(a-b) - \cos(a+b)}{2},$$

$$\cos(a)\sin(b) = \frac{\sin(a+b) - \sin(a-b)}{2}.$$

If a and b are integers then $a \pm b$ are integers. When $p \neq 0$ is a integer

$$\int_{-\pi}^{\pi} \cos(px) dx = 0, \quad \int_{-\pi}^{\pi} \sin(px) dx = 0.$$

When $f \neq g$ and when $f = g$

If f and g are from the list

$$1, \cos(x), \sin(x), \cos(2x), \sin(2x), \dots, \cos(nx), \sin(nx), \dots$$

and $f \neq g$ then

$$\int_{-\pi}^{\pi} f(x)g(x) dx = 0.$$

When $f(x) = g(x)$ we have the following cases.

$$\int_{-\pi}^{\pi} dx = 2\pi,$$

$$\int_{-\pi}^{\pi} \cos^2(nx) dx = \pi,$$

$$\int_{-\pi}^{\pi} \sin^2(nx) dx = \pi.$$

The formula for the Fourier coefficients a_n and b_n

If our starting point are numbers a_n and b_n such that the series converges to define

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(nx) + b_n \sin(nx))$$

and the convergence is such that all operations such as interchanging infinite sums and integrals are valid then the orthogonality properties imply the following.

$$\begin{aligned} \int_{-\pi}^{\pi} f(x) dx &= a_0\pi, \\ \int_{-\pi}^{\pi} f(x) \cos(mx) dx &= a_m\pi, \\ \int_{-\pi}^{\pi} f(x) \sin(mx) dx &= b_m\pi. \end{aligned}$$

This gives a justification for the formulas used.

The Heaviside function on $(-\pi, \pi]$

Let

$$f_1(x) = \begin{cases} 1, & \text{if } 0 \leq x \leq \pi, \\ 0, & \text{if } -\pi < x < 0 \end{cases}$$

which we continue in a 2π -periodic way.


Determining the coefficients gives the following.

$$f_1(x) \sim \frac{1}{2} + \frac{2}{\pi} \left(\sin(x) + \frac{\sin(3x)}{3} + \dots + \frac{\sin((2n-1)x)}{2n-1} + \dots \right).$$

We can immediately observe that when $x = k\pi$ the value of the series is $1/2$. The series does not converge to $f_1(x)$ at these points of discontinuity.

A plot of a truncated Fourier series for $f_1(x)$

$$S_{63}(x) = \frac{1}{2} + \frac{2}{\pi} \left(\sin(x) + \frac{\sin(3x)}{3} + \dots + \frac{\sin(63x)}{63} \right).$$



The function $f_2(x) = |x|$ on $(-\pi, \pi]$

In this case convergence occurs at all points and we can write for $|x| \leq \pi$.

$$|x| = \frac{\pi}{2} - \frac{4}{\pi} \left(\cos(x) + \frac{\cos(3x)}{3^2} + \dots + \frac{\cos((2n-1)x)}{(2n-1)^2} + \dots \right).$$


The 2π -periodic function extension is continuous at all points.

Observe that $f_2(x)$ is an even function of x which is why $b_n = 0$ and only cosine terms are involved. Integration by parts is used to determine

$$a_n = \frac{2}{\pi} \int_0^\pi x \cos(nx) dx.$$

A plot of a truncated Fourier series for $f_2(x)$

$$S_7(x) = \frac{\pi}{2} - \frac{4}{\pi} \left(\cos(x) + \frac{\cos(3x)}{3^2} + \frac{\cos(5x)}{5^2} + \frac{\cos(7x)}{7^2} \right).$$



The function $f_3(x) = x$ on $(-\pi, \pi]$

In this case for $x \in (-\pi, \pi)$ we have convergence to $f_3(x)$ and we can write

$$x = 2 \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin(nx) = 2 \left(\sin(x) - \frac{\sin(2x)}{2} + \frac{\sin(3x)}{3} - \dots \right).$$

We can immediately observe that when $x = \pi$ the value of the series is 0. The series does converge here but not to $f_3(\pi) = \pi$ or to the limit as we tend to $-\pi$ which is $-\pi$. The 2π -periodic extension is discontinuous at the points $(2k + 1)\pi$, $k \in \mathbb{Z}$.

Observe that $f_3(x)$ is an odd function of x which is why $a_n = 0$ and only sine terms are involved. Integration by parts is used to determine

$$b_n = \frac{2}{\pi} \int_0^{\pi} x \sin(nx) dx.$$

A plot of a truncated Fourier series for $f_3(x)$

$$S_{64}(x) = 2 \sum_{n=1}^{64} \frac{(-1)^{n+1}}{n} \sin(nx).$$

Fourier series for f3 with terms up to sin(64x)

