

## MA3614 Complex variable methods and applications Comments, topics and why it is taught

### ▶ Will the module involve complex numbers?

Yes. The complex number material that you learned in MA1620 will be used.

The module is more about functions of a complex variable. For many real valued functions  $f(x)$ ,  $x \in \mathbb{R}$  it makes sense to consider

$$f(z), \quad z = x + iy, \quad x, y \in \mathbb{R}, \quad i^2 = -1.$$

The natural domain of many functions that you have considered is the complex plane. Hence you learn more about such functions.

MA3614 Welcome Slides for 2023/4 Page 1 of 12

### What previous study will be useful?

- ▶ Complex number manipulation from MA1620, e.g.  $z = x + iy = re^{i\theta}$ ,  $z^n = r^n e^{in\theta}$  etc.
- ▶ Partial differentiation from MA2612, e.g. for a sufficiently smooth function  $u(x, y)$ ,

$$\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x}.$$

- ▶ Geometric series from possibly several previous modules, i.e.

$$\frac{1}{1-z} = 1 + z + z^2 + \cdots + z^n + \cdots, \quad \text{when } |z| < 1.$$

MA3614 Welcome Slides for 2023/4 Page 3 of 12

## Comments, topics and why it is taught continued

### ▶ Why study something that is not real?

A brief answer to this is that it helps to understand the real case better. There are some examples of this in these slides.

It is also a tool in solving real problems. This is the application part.

MA3614 Welcome Slides for 2023/4 Page 2 of 12

### Without detail what topics are involved?

- ▶ Differentiation in a complex sense.
- ▶ Integration in the complex plane.
- ▶ Power series and Laurent series representations of functions. (Term 2).
- ▶ Applications usually involving residue theory. (Term 2).

MA3614 Welcome Slides for 2023/4 Page 4 of 12

# Which functions make sense with a complex variable?

## 1. Polynomials

$$p(z) = a_0 + a_1z + \cdots + a_nz^n, \quad a_n \neq 0.$$

This has  $n$  roots (counting multiplicities) in the complex plane. We need to study complex integration to explain this.

## 2. Rational functions (i.e. a ratio of polynomials).

$$f(z) = \frac{a_0 + a_1z + \cdots + a_nz^n}{b_0 + b_1z + \cdots + b_mz^m}.$$

When  $n < m$  there is a partial fraction representation. You may have had rules to get this representation. Do you know why the rules work?

## 3. Exponential function.

$$\exp(z) = e^x e^{iy} = e^x (\cos(y) + i \sin(y)).$$

The real case of  $e^x$  and the notation  $e^{i\theta} = \cos(\theta) + i \sin(\theta)$  are special cases.

## What additional properties will be covered? Differentiation and analytic

In the real case differentiation is considered. In this module we consider when the functions are also differentiable in a complex sense and a related **analytic** property. Many additional results will depend on where  $f(z)$  is analytic and where it is not. The complex differentiable property at  $z_0$  is concerned with when the following limit exists.

$$\frac{df}{dz}(z_0) \equiv f'(z_0) := \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}.$$

We have the same expression as in the real case but now we are dividing by a complex number and we must get the same value however  $h$  tends to 0 to be complex differentiable at  $z_0$ .

$f(z)$  is analytic at  $z_0$  if it is complex differentiable at  $z_0$  and in a neighbourhood of  $z_0$ . This will probably first be done in about week 3.

# Some examples of things the complex case explains

The following relate to things you possibly have met before.

## 1. Suppose that you have a real polynomial.

$$p(x) = a_0 + a_1x + \cdots + a_nx^n, \quad a_n \neq 0, \quad a_k \in \mathbb{R}.$$

Non-real roots occur in complex conjugate pairs. This is a consequence of

$$p(\bar{z}) = \overline{p(z)}.$$

## 2. Why is the Maclaurin series for

$$f(x) = \frac{1}{1+x^2} = 1 + (-x^2) + (-x^2)^2 + (-x^2)^3 + (-x^2)^4 + \cdots$$

only valid in  $-1 < x < 1$ ? Note that the function is infinitely differentiable on  $\mathbb{R}$ . This is because  $f(z)$  has singularities at  $\pm i$ . The series (which is a geometric series) is valid for  $|z| < 1$ .

## Why is MA2612 a prerequisite?

With  $z = x + iy$ ,  $x, y \in \mathbb{R}$  a function of a complex variable

$$w = f(z), \quad w = u + iv, \quad u, v \in \mathbb{R}$$

is in full

$$f(x + iy) = u(x, y) + iv(x, y).$$

We have real valued functions  $u$  and  $v$  of 2-variables  $x, y$ . When  $f(z)$  is complex differentiable we can express  $f'(z)$  in terms of the partial derivatives

$$\frac{\partial u}{\partial x}, \quad \frac{\partial v}{\partial x}, \quad \frac{\partial u}{\partial y} \quad \text{and} \quad \frac{\partial v}{\partial y}.$$

We will see that  $f(z)$  is analytic in a domain if and only if the following hold in the domain.

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

These are the Cauchy Riemann equations.

## Contour integrals

In the real case you consider definite integrals of the form

$$\int_a^b f(x) dx.$$


Generalising to the complex case involves an arc  $\Gamma$  in the complex plane and we write

$$\int_{\Gamma} f(z) dz.$$

### Examples of $\Gamma$



A line segment.



A union of a line segment and a half circle to give a loop.

MA3614 Welcome Slides for 2023/4 Page 9 of 12

## Early jargon: Laurent series

A Laurent series is a series of the form

$$\sum_{n=-\infty}^{\infty} a_n (z - z_0)^n.$$

When it converges the region is an annulus  $\{z : r < |z - z_0| < R\}$ .

## Laurent series representation

Let  $f(z)$  be analytic in an annulus  $r < |z - z_0| < R$ . Then it has the representation

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} a_{-n} (z - z_0)^{-n}.$$

MA3614 Welcome Slides for 2023/4 Page 11 of 12

## Taylor series will be explained?

With integration introduced a key result early in term 2 is to show that when  $f(z)$  is analytic in a domain,  $\Gamma$  is a closed loop traversed once in the anti-clockwise direction and  $z$  is inside  $\Gamma$  we have the Cauchy integral formula

$$f(z) = \frac{1}{2\pi i} \oint_{\Gamma} \frac{f(\zeta)}{\zeta - z} d\zeta.$$

This implies the generalised Cauchy integral formula

$$f^{(n)}(z) = \frac{n!}{2\pi i} \oint_{\Gamma} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta, \quad n = 0, 1, 2, \dots$$

Using both gives the Taylor series

$$f(z) = \sum_{k=0}^{\infty} \frac{f^{(k)}(z_0)}{k!} (z - z_0)^k.$$

If  $f(z)$  is analytic in  $|z - z_0| < R$  then the series representation is valid in this disk.

MA3614 Welcome Slides for 2023/4 Page 10 of 12

## Early jargon: A residue

This will first be met when considering partial fractions. Consider

$$R(z) = \frac{p(z)}{q(z)}, \quad q(z) = (z - z_1)(z - z_2) \cdots (z - z_n).$$

When  $\deg p(z) < \deg q(z)$  and the zeros of  $q(z)$  are simple we have the partial fraction representation of the form

$$R(z) = \frac{p(z)}{q(z)} = \sum_{k=1}^n \frac{A_k}{z - z_k}.$$

Here  $A_k$  is the **residue** at  $z_k$ . This will be covered in term 1.

More generally, when

$$\sum_{n=-\infty}^{\infty} a_n (z - z_0)^n$$

converges in  $0 < |z - z_0| < R$  the coefficient  $a_{-1}$  is the residue at  $z_0$ . This will be covered in term 2.

MA3614 Welcome Slides for 2023/4 Page 12 of 12